

Product datasheet for **TA378157**

Lipolysis Stimulated Lipoprotein Receptor (LSR) Rabbit Polyclonal Antibody


Product data:

| | |
|-------------------------|---|
| Product Type: | Primary Antibodies |
| Applications: | WB |
| Recommended Dilution: | WB,1:500 - 1:2000 |
| Reactivity: | Human, Mouse, Rat |
| Modifications: | Unmodified |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 50-240 of human LSR (NP_057009.3). |
| Formulation: | Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| Concentration: | lot specific |
| Purification: | Affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. |
| Stability: | Shelf life: one year from despatch. |
| Predicted Protein Size: | 54kDa/64kDa/65kDa/69kDa/71kDa |
| Gene Name: | lipolysis stimulated lipoprotein receptor |
| Database Link: | Entrez Gene 51599 Human Q86X29 |
| Background: | Probable role in the clearance of triglyceride-rich lipoprotein from blood. Binds chylomicrons, LDL and VLDL in presence of free fatty acids and allows their subsequent uptake in the cells (By similarity). |
| Synonyms: | ILDR3; LISCH; LISCH7; MGC10659; MGC48312; MGC48503 |



[View online »](#)

Product images:



Western blot analysis of extracts of various cell lines, using LSR antibody (TA378157) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 10s.