

## Product datasheet for **TA378365**

### MAX Rabbit Polyclonal Antibody

#### Product data:


Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Human
Modifications:	Phospho S11
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	A synthetic phosphorylated peptide around S11 of human MAX (NP_002373.3).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	9kDa/11kDa/12kDa/15kDa/17kDa/18kDa
Gene Name:	MYC associated factor X
Database Link:	<a href="#">Entrez Gene 4149 Human P61244</a>
Background:	The protein encoded by this gene is a member of the basic helix-loop-helix leucine zipper (bHLHZ) family of transcription factors. It is able to form homodimers and heterodimers with other family members, which include Mad, Mxi1 and Myc. Myc is an oncoprotein implicated in cell proliferation, differentiation and apoptosis. The homodimers and heterodimers compete for a common DNA target site (the E box) and rearrangement among these dimer forms provides a complex system of transcriptional regulation. Mutations of this gene have been reported to be associated with hereditary pheochromocytoma. A pseudogene of this gene is located on the long arm of chromosome 7. Alternative splicing results in multiple transcript variants.



[View online »](#)

Synonyms: bHLHd4; bHLHd5; bHLHd6; bHLHd7; bHLHd8; MGC10775; MGC11225; MGC18164; MGC34679; MGC36767; orf1

### Product images:



Western blot analysis of extracts of Jurkat cells, using Phospho-MAX-S11 antibody (TA378365) at 1:1000 dilution. Jurkat cells were treated by serum-starvation overnight. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA.