

## **Product datasheet for TA378401S**

## Product data:

**Product Type:** Primary Antibodies

MCM2 Rabbit Polyclonal Antibody

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human

Modifications: Phospho S139

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthetic phosphorylated peptide around S139 of human MCM2.

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 101kDa

**Gene Name:** minichromosome maintenance complex component 2

Database Link: Entrez Gene 4171 Human

P49736

**Background:** The protein encoded by this gene is one of the highly conserved mini-chromosome

maintenance proteins (MCM) that are involved in the initiation of eukaryotic genome replication. The hexameric protein complex formed by MCM proteins is a key component of

the pre-replication complex (pre\_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. This protein forms a complex with MCM4, 6, and 7, and has been shown to regulate the helicase activity of the complex. This protein is phosphorylated, and thus regulated by, protein kinases CDC2 and CDC7. Multiple alternatively spliced transcript variants have been found, but the full-length

nature of some variants has not been defined.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

BM28; CCNL1; cdc19; CDCL1; D3S3194; KIAA0030; MGC10606; MITOTIN

## **Product images:**



Western blot analysis of extracts of HeLa cells, using Phospho-MCM2-S139 antibody ([TA378401]) at 1:1000 dilution.Hela cells were treated by CIP(20uL/400ul) at 37°C for 1 hour.HeLa cells were treated by Serum-starvation overnight at 37°C.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit.|Exposure time: 180s.