

Product datasheet for TA378543

i Toddet datasiieet for TAS7054

Product data:

Product Type: Primary Antibodies

MRTFA Rabbit Polyclonal Antibody

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:100 IF,1:50 - 1:100

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of

human MKL1 (NP_065882.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 98kDa

Gene Name: megakaryoblastic leukemia (translocation) 1

Database Link: Entrez Gene 57591 Human

Q969V6

Background: The protein encoded by this gene interacts with the transcription factor myocardin, a key

regulator of smooth muscle cell differentiation. The encoded protein is predominantly nuclear and may help transduce signals from the cytoskeleton to the nucleus. This gene is involved in a specific translocation event that creates a fusion of this gene and the RNA-binding motif protein-15 gene. This translocation has been associated with acute

megakaryocytic leukemia. Alternative splicing results in multiple transcript variants.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

AMKL; BSAC; KIAA1438; MAL; MRTF-A; OTTHUMP00000199247

Product images:



Western blot analysis of extracts of various cell lines, using MKL1 antibody (TA378543) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.



Immunofluorescence analysis of U-2 OS cells using MKL1 Polyclonal Antibody (TA378543) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.