

Product datasheet for **TA378566**

AF9 (MLLT3) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-150 of human MLLT3/AF9 (NP_004520.2).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	63kDa
Gene Name:	myeloid/lymphoid or mixed-lineage leukemia; translocated to, 3
Database Link:	Entrez Gene 4300 Human P42568




[View online »](#)

Background:

Chromatin reader component of the super elongation complex (SEC, a complex required to increase the catalytic rate of RNA polymerase II transcription by suppressing transient pausing by the polymerase at multiple sites along the DNA. Specifically recognizes and binds acylated histone H3, with a preference for histone H3 that is crotonylated. Crotonylation marks active promoters and enhancers and confers resistance to transcriptional repressors. Recognizes and binds histone H3 crotonylated at 'Lys-9' (H3K9cr, and with slightly lower affinity histone H3 crotonylated at 'Lys-18' (H3K18cr. Also recognizes and binds histone H3 acetylated and butyrylated at 'Lys-9' (H3K9ac and H3K9bu, respectively, but with lower affinity than crotonylated histone H3. In the SEC complex, MLLT3 is required to recruit the complex to crotonylated histones. Recruitment of the SEC complex to crotonylated histones promotes recruitment of DOT1L on active chromatin to deposit histone H3 'Lys-79' methylation (H3K79me. Plays a key role in hematopoietic stem cell (HSC maintenance by preserving, rather than conferring, HSC stemness. Acts by binding to the transcription start site of active genes in HSCs and sustaining level of H3K79me2, probably by recruiting DOT1L.

Synonyms:

AF-9; AF9; FLJ2035; YEATS3

Product images:

Western blot analysis of extracts of various cell lines, using MLLT3/AF9 antibody (TA378566) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 30s.