

## Product datasheet for **TA378607**

### MOV10 Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:1000 - 1:2000
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-310 of human MOV10 (NP_066014.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	6kDa/101kDa/113kDa
Gene Name:	Mov10 RISC complex RNA helicase
Database Link:	<a href="#">Entrez Gene 4343 Human Q9HCE1</a>




[View online »](#)

**Background:**

5' to 3' RNA helicase contributing to UPF1 mRNA target degradation by translocation along 3' UTRs. Required for microRNA (miRNA-mediated gene silencing by the RNA-induced silencing complex (RISC. Required for both miRNA-mediated translational repression and miRNA-mediated cleavage of complementary mRNAs by RISC. In cooperation with FMR1, regulates miRNA-mediated translational repression by AGO2. Restricts retrotransposition of long interspersed element-1 (LINE-1 in cooperation with TUT4 and TUT7 counteracting the RNA chaperone activity of L1RE1. Facilitates LINE-1 uridylation by TUT4 and TUT7. Required for embryonic viability and for normal central nervous system development and function. Plays two critical roles in early brain development: suppresses retroelements in the nucleus by directly inhibiting cDNA synthesis, while regulates cytoskeletal mRNAs to influence neurite outgrowth in the cytosol (By similarity. May function as a messenger ribonucleoprotein (mRNP clearance factor. Exhibits antiviral activity against dengue virus (DENV.

**Synonyms:**

DKFZp667O1423; FLJ32791; fSAP113; gb110; KIAA1631; MGC2948; OTTHUMP00000013679; OTTHUMP00000013680

**Product images:**

Western blot analysis of extracts of various cell lines, using MOV10 antibody (TA378607) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 30s.