

### **Product datasheet for TA378723**

# **ATP6 Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:100 - 1:200

IF,1:50 - 1:200

Reactivity: Human, Mouse

Modifications: Unmodified

**Host:** Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic peptide corresponding to a sequence within amino acids 100 to the C-terminus of

mouse ATP6 (NP\_904333.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 25kDa

**Gene Name:** ATP synthase 6, mitochondrial

**Database Link:** Entrez Gene 17705 Mouse

P00848



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

Mitochondrial membrane ATP synthase (F(1F(0 ATP synthase or Complex V produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1 - containing the extramembraneous catalytic core and F(0 - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1 is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Key component of the proton channel; it may play a direct role in the translocation of protons across the membrane.

## **Product images:**



Western blot analysis of extracts of various cell lines, using ATP6 antibody (TA378723) at 1:1000 dilution.\_Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.\_Lysates/proteins: 25ug per lane.\_Blocking buffer: 3% nonfat dry milk in TBST.\_Detection: ECL Enhanced Kit .\_Exposure time: 10s.



Immunofluorescence analysis of U2OS cells using ATP6 antibody (TA378723) at dilution of 1:100. Blue: DAPI for nuclear staining.