

Product datasheet for TA378803

MUS81 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:100 IF,1:50 - 1:100

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-350 of

human MUS81 (NP_079404.3).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 61kDa

Gene Name: MUS81 structure-specific endonuclease subunit

Database Link: Entrez Gene 80198 Human

Q96NY9



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

This gene encodes a structure-specific endonuclease which belongs to the XPF/MUS81 endonuclease family and plays a critical role in the resolution of recombination intermediates during DNA repair after inter-strand cross-links, replication fork collapse, and DNA double-strand breaks. The encoded protein associates with one of two closely related essential meiotic endonuclease proteins (EME1 or EME2) to form a complex that processes DNA secondary structures. It contains an N-terminal DEAH helicase domain, an excision repair cross complementation group 4 (ERCC4) endonuclease domain, and two tandem C-terminal helix-hairpin-helix domains. Mice with a homozygous knockout of the orthologous gene have significant meiotic defects including the failure to repair a subset of DNA double strand breaks.

Synonyms: FLJ21012; FLJ44872

Product images:



Western blot analysis of extracts of various cell lines, using MUS81 antibody (TA378803) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit. |Exposure time: 90s.