

## **Product datasheet for TA378816**

### c-Myc (MYC) Rabbit Polyclonal Antibody

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** ELISA, WB

Recommended Dilution: WB.1:500 - 1:1000

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse

**Modifications:** Unmodified

Rabbit Host:

Isotype: **IgG** 

Clonality: Polyclonal

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles. Storage:

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** 51kDa

Gene Name: v-myc avian myelocytomatosis viral oncogene homolog

Database Link: Entrez Gene 4609 Human

P01106



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:

This gene is a proto-oncogene and encodes a nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. The encoded protein forms a heterodimer with the related transcription factor MAX. This complex binds to the E box DNA consensus sequence and regulates the transcription of specific target genes. Amplification of this gene is frequently observed in numerous human cancers. Translocations involving this gene are associated with Burkitt lymphoma and multiple myeloma in human patients. There is evidence to show that translation initiates both from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site, resulting in the production of two isoforms with distinct N-termini.

**Synonyms:** bHLHe39; c-Myc; MRTL

# **Product images:**



Western blot analysis of lysates from HeLa cells