

## **Product datasheet for TA378917**

## **NAIP Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:2000

**Reactivity:** Rat

Host: Rabbit

**Isotype:** IgG

Clonality: Polyclonal

**Immunogen:** A synthetic peptide corresponding to a sequence within amino acids 750-850 of human NAIP

(NP\_075043.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** NLR family, apoptosis inhibitory protein

Database Link: Q13075

**Background:** This gene is part of a 500 kb inverted duplication on chromosome 5q13. This duplicated

region contains at least four genes and repetitive elements which make it prone to

rearrangements and deletions. The repetitiveness and complexity of the sequence have also caused difficulty in determining the organization of this genomic region. This copy of the gene is full length; additional copies with truncations and internal deletions are also present in this region of chromosome 5q13. It is thought that this gene is a modifier of spinal muscular atrophy caused by mutations in a neighboring gene, SMN1. The protein encoded by this gene

contains regions of homology to two baculovirus inhibitor of apoptosis proteins, and it is able

to suppress apoptosis induced by various signals. Alternative splicing and the use of

alternative promoters results in multiple transcript variants.

Synonyms: BIRC1; FLJ18088; FLJ42520; FLJ58811; NLRB1; OTTHUMP00000125255; psiNAIP




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Western blot analysis of extracts of various cell lines, using NAIP antibody (TA378917) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit. |Exposure time: 10s.