

#### **Product datasheet for TA378917**

# **OriGene Technologies, Inc.**9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **NAIP Rabbit Polyclonal Antibody**

#### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ELISA, WB

Recommended Dilution: WB,1:500 - 1:2000

ELISA,Recommended starting concentration is 1 μg/mL. Please optimize the concentration

based on your specific assay requirements.

**Reactivity:** Rat

**Host:** Rabbit

**Isotype:** IgG

**Clonality:** Polyclonal

**Formulation:** Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Concentration: lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 160kDa

**Gene Name:** NLR family, apoptosis inhibitory protein

Database Link: Q13075

**Background:** This gene is part of a 500 kb inverted duplication on chromosome 5q13. This duplicated

region contains at least four genes and repetitive elements which make it prone to

rearrangements and deletions. The repetitiveness and complexity of the sequence have also caused difficulty in determining the organization of this genomic region. This copy of the gene is full length; additional copies with truncations and internal deletions are also present in this region of chromosome 5q13. It is thought that this gene is a modifier of spinal muscular atrophy caused by mutations in a neighboring gene, SMN1. The protein encoded by this gene contains regions of homology to two baculovirus inhibitor of apoptosis proteins, and it is able

to suppress apoptosis induced by various signals. Alternative splicing and the use of

alternative promoters results in multiple transcript variants.






Synonyms:

BIRC1; FLJ18088; FLJ42520; FLJ58811; NLRB1; OTTHUMP00000125255; psiNAIP

## **Product images:**



Western blot analysis of various lysates using NAIP Rabbit pAb (TA378917) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Blocking buffer: 3% nonfat dry milk in TBST Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.