

Product datasheet for TA378985

Product data:

Product Type: Primary Antibodies

NDRG1 Rabbit Polyclonal Antibody

Applications: WB

Reactivity: WB,1:500 - 1:2000

Reactivity: Human, Mouse

Modifications: Phospho S330

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic phosphorylated peptide around S330 of human NDRG1 (NP_001128714.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 33kDa/35kDa/42kDa

Gene Name: N-myc downstream regulated 1

Database Link: Entrez Gene 10397 Human

Q92597

Background: This gene is a member of the N-myc downregulated gene family which belongs to the

alpha/beta hydrolase superfamily. The protein encoded by this gene is a cytoplasmic protein involved in stress responses, hormone responses, cell growth, and differentiation. The encoded protein is necessary for p53-mediated caspase activation and apoptosis. Mutations in this gene are a cause of Charcot-Marie-Tooth disease type 4D, and expression of this gene may be a prognostic indicator for several types of cancer. Alternatively spliced transcript

variants encoding multiple isoforms have been observed for this gene.

Synonyms: CAP43; CMT4D; DRG-1; DRG1; GC4; HMSNL; NDR1; NMSL; PROXY1; RIT42; RTP; TARG1; TDD5




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of extracts of Jurkat cells, using Phospho-NDRG1-S330 pAb (TA378985) at 1:2000 dilution or NDRG1 antibody (A2142). Jurkat cells were treated by Calyculin A (100nM) for 30 minutes. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA. | Detection: ECL Basic Kit. | Exposure time: 1s.