

Product datasheet for TA379068

Nexilin (NEXN) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 516-675 of

human NEXN (NP_653174.3).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 72kDa/73kDa/79kDa/80kDa

Gene Name: nexilin F-actin binding protein

Database Link: Entrez Gene 91624 Human

Q0ZGT2

Background: This gene encodes a filamentous actin-binding protein that may function in cell adhesion and

migration. Mutations in this gene have been associated with dilated cardiomyopathy, also

known as CMD1CC. Alternatively spliced transcript variants have been described.

Synonyms: CMD1CC; MGC104234; MGC138865; MGC138866; NELIN; nexilin




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of extracts of various cell lines, using NEXN antibody (TA379068) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.



Immunofluorescence analysis of U-2 OS cells using NEXN antibody (TA379068) at dilution of 1:100. Blue: DAPI for nuclear staining.