

Product datasheet for TA379117

NGF Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:1000

Reactivity: Mouse, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 19-241 of

human NGF (NP_002497.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 26kDa

Gene Name: nerve growth factor

Database Link: P01138

Background: This gene is a member of the NGF-beta family and encodes a secreted protein which

homodimerizes and is incorporated into a larger complex. This protein has nerve growth stimulating activity and the complex is involved in the regulation of growth and the

differentiation of sympathetic and certain sensory neurons. Mutations in this gene have been

associated with hereditary sensory and autonomic neuropathy, type 5 (HSAN5), and

dysregulation of this gene's expression is associated with allergic rhinitis.

Synonyms: Beta-NGF; HSAN5; MGC161426; MGC161428; NGFB; NID67




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:





Western blot analysis of extracts of Mouse liver, using NGF antibody (TA379117) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 180s.

Western blot analysis of extracts of Rat liver, using NGF antibody (TA379117) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 180s.