

## **Product datasheet for TA379183**

## **NMNAT3 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 1-215 of

human NMNAT3 (NP\_835471.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 18kDa/24kDa/28kDa

**Gene Name:** nicotinamide nucleotide adenylyltransferase 3

Database Link: Entrez Gene 349565 Human

Q96T66

**Background:** This gene encodes a member of the nicotinamide/nicotinic acid mononucleotide

adenylyltransferase family. These enzymes use ATP to catalyze the synthesis of nicotinamide

adenine dinucleotide or nicotinic acid adenine dinucleotide from nicotinamide

mononucleotide or nicotinic acid mononucleotide, respectively. The encoded protein is localized to mitochondria and may also play a neuroprotective role as a molecular

chaperone. Alternatively spliced transcript variants encoding multiple isoforms have been

observed for this gene.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




**Synonyms:** PNAT-3; PNAT3

## **Product images:**




Western blot analysis of extracts of various cell lines, using NMNAT3 antibody (TA379183) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.



Immunofluorescence analysis of HeLa cells using NMNAT3 Rabbit pAb (TA379183) at dilution of 1:100. Blue: DAPI for nuclear staining.





Immunofluorescence analysis of NIH/3T3 cells using NMNAT3 Rabbit pAb (TA379183) at dilution of 1:100. Blue: DAPI for nuclear staining.