

## **Product datasheet for TA379223S**

## **eNOS (NOS3) Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200

**Reactivity:** Human, Mouse **Modifications:** Phospho S1177

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic phosphorylated peptide around S1177 of human eNOS (NP\_000594.2).

**Formulation:** PBS with 0.05% proclin300,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 67kDa/68kDa/133kDa

Gene Name: nitric oxide synthase 3

Database Link: Entrez Gene 4846 Human

P29474

**Background:** Nitric oxide is a reactive free radical which acts as a biologic mediator in several processes,

including neurotransmission and antimicrobial and antitumoral activities. Nitric oxide is synthesized from L-arginine by nitric oxide synthases. Variations in this gene are associated

with susceptibility to coronary spasm. Alternative splicing and the use of alternative

promoters results in multiple transcript variants.

**Synonyms:** cNOS; EC-NOS; ECNOS; eNOS; NOSIII




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Western blot analysis of extracts of SKOV3 cells, using Phospho-eNOS-S1177 antibody ([TA379223]). SKOV3 cells were treated by EGF. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA.



Western blot analysis of extracts of NIH/3T3 cells, using Phospho-eNOS-S1177 antibody ([TA379223]) at 1:1000 dilution. NIH/3T3 cells were treated by EGF (100ng/mL) for 30 minutes after serum-starvation overnight. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA. | Detection: ECL Enhanced Kit. | Exposure time: 60s.