

## **Product datasheet for TA379281S**

# **NR0B2** Rabbit Polyclonal Antibody

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** ELISA, WB

Recommended Dilution: WB.1:500 - 1:1000

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Mouse, Rat **Modifications:** Unmodified

Rabbit Host: Isotype: **IgG** 

Clonality: Polyclonal

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles. Storage:

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** 28kDa

Gene Name: nuclear receptor subfamily 0 group B member 2

Database Link: Q15466



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




### Background:

The protein encoded by this gene is an unusual orphan receptor that contains a putative ligand-binding domain but lacks a conventional DNA-binding domain. The gene product is a member of the nuclear hormone receptor family, a group of transcription factors regulated by small hydrophobic hormones, a subset of which do not have known ligands and are referred to as orphan nuclear receptors. The protein has been shown to interact with retinoid and thyroid hormone receptors, inhibiting their ligand-dependent transcriptional activation. In addition, interaction with estrogen receptors has been demonstrated, leading to inhibition of function. Studies suggest that the protein represses nuclear hormone receptor-mediated transactivation via two separate steps: competition with coactivators and the direct effects of its transcriptional repressor function.

**Synonyms:** FLJ17090; SHP; SHP1

# **Product images:**



Western blot analysis of various lysates using SHP/NR0B2 Rabbit pAb ([TA379281]) at 1:900 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.