

Product datasheet for TA379302

Nurr1 (NR4A2) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: WB,1:500 - 1:2000 Human, Mouse

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic Peptide of human NR4A2.

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 59kDa/66kDa

Gene Name: nuclear receptor subfamily 4 group A member 2

Database Link: Entrez Gene 4929 Human

P43354

Background: This gene encodes a member of the steroid-thyroid hormone-retinoid receptor superfamily.

The encoded protein may act as a transcription factor. Mutations in this gene have been associated with disorders related to dopaminergic dysfunction, including Parkinson disease, schizophernia, and manic depression. Misregulation of this gene may be associated with rheumatoid arthritis. Alternatively spliced transcript variants have been described, but their

biological validity has not been determined.

Synonyms: HZF-3; NOT; NURR1; OTTHUMP00000204707; RNR1; TINUR




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western blot analysis of extracts of various cell lines, using (TA379302) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 180s.



Western blot analysis of extracts of Mouse brain, using (TA379302) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 180s.