

Product datasheet for **TA379308**


NRBF2 Rabbit Polyclonal Antibody

Product data:


| | |
|-------------------------|---|
| Product Type: | Primary Antibodies |
| Applications: | IF, IHC, WB |
| Recommended Dilution: | WB,1:500 - 1:2000 IHC,1:50 - 1:200 IF,1:50 - 1:200 |
| Reactivity: | Human, Mouse, Rat |
| Modifications: | Unmodified |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 10-220 of human NRBF2 (NP_110386.2). |
| Formulation: | Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| Concentration: | lot specific |
| Purification: | Affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. |
| Stability: | Shelf life: one year from despatch. |
| Predicted Protein Size: | 26kDa/31kDa/32kDa |
| Gene Name: | nuclear receptor binding factor 2 |
| Database Link: | Entrez Gene 29982 Human Q96F24 |
| Background: | May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro). |
| Synonyms: | COPR; COPR1; COPR2; DKFZp564C1664; FLJ30395; NRBF-2 |




[View online »](#)

Product images:


Western blot analysis of extracts from normal (control) and NRBF2 knockout (KO) 293T cells, using NRBF2 antibody (TA379308) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 10s.



Western blot analysis of extracts of various cell lines, using NRBF2 antibody (TA379308) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit . | Exposure time: 90s.



Immunohistochemistry of paraffin-embedded human esophageal using NRBF2 Rabbit pAb (TA379308) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.