

## **Product datasheet for TA379313S**

## **NRG4 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 1-115 of

human NRG4 (NP\_612640.1).

**Formulation:** PBS with 0.05% proclin300,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 12kDa

**Gene Name:** neuregulin 4

Database Link: Entrez Gene 145957 Human

Q8WWG1

Background: The neuregulins, including NRG4, activate type-1 growth factor receptors (see EGFR; MIM

131550) to initiating cell-to-cell signaling through tyrosine phosphorylation (Harari et al., 1999

[PubMed 10348342]).[supplied by OMIM, Mar 2008]

Synonyms: DKFZp779N0541; DKFZp779N1944; HRG4; Pro-NRG4




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Western blot analysis of extracts of various cell lines, using NRG4 antibody ([TA379313]) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Enhanced Kit. |Exposure time: 3min.