

Product datasheet for TA379490S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OR2T10 Rabbit Polyclonal Antibody

Product data:

Isotype:

Product Type: Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: WB.1:500 - 1:2000

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse

Modifications: Unmodified

Host: Rabbit **IgG**

Clonality: Polyclonal

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles. Storage:

Shelf life: one year from despatch. Stability:

Predicted Protein Size: 35kDa

Gene Name: olfactory receptor family 2 subfamily T member 10

Database Link: Entrez Gene 127069 Human

Q8NGZ9

Background: Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal

response that triggers the perception of a smell. The olfactory receptor proteins are

members of a large family of G-protein-coupled receptors (GPCR) arising from single codingexon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and


proteins for this organism is independent of other organisms.





Synonyms: OR1-64

Product images:



Western blot analysis of various lysates using OR2T10 Rabbit pAb ([TA379490]) at 1:1000 dilution.

Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.