

Product datasheet for TA379577

LIS1 (PAFAH1B1) Rabbit Polyclonal Antibody

Product data:

Reactivity:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB,1:500 - 1:2000

Modifications: Unmodified

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-170 of

human PAFAH1B1 (NP_000421.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Human, Mouse

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 16kDa/46kDa

Gene Name: platelet activating factor acetylhydrolase 1b regulatory subunit 1

Database Link: Entrez Gene 5048 Human

P43034

Background: This locus was identified as encoding a gene that when mutated or lost caused the

lissencephaly associated with Miller-Dieker lissencephaly syndrome. This gene encodes the

non-catalytic alpha subunit of the intracellular lb isoform of platelet-activating factor

acteylhydrolase, a heterotrimeric enzyme that specifically catalyzes the removal of the acetyl group at the SN-2 position of platelet-activating factor (identified as 1-O-alkyl-2-acetyl-sn-glyceryl-3-phosphorylcholine). Two other isoforms of intracellular platelet-activating factor acetylhydrolase exist: one composed of multiple subunits, the other, a single subunit. In

addition, a single-subunit isoform of this enzyme is found in serum.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

LIS-1; LIS1; LIS2; MDCR; MDS; PAFAH; PAFAHA

Product images:



Western blot analysis of extracts of various cell lines, using PAFAH1B1 antibody (TA379577) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 10s.