

## Product datasheet for **TA379599**

### Mono-Methyl lysine Rabbit Polyclonal Antibody


#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000 IP,1:50 - 1:200 ChIP,1:50 - 1:200
Reactivity:	Human, Mouse, Rat
Modifications:	pan-Mono-Methyl K
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	A synthetic peptide corresponding to a sequence containing monomethylated K.
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.




[View online »](#)


**Product images:**



Western blot analysis of extracts of various cell lines, using Pan Mono-Methyl lysine antibody (TA379599) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit . |Exposure time: 10s.



Western blot analysis of extracts of various cell lines, using Pan Mono-Methyl lysine Rabbit pAb (TA379599) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Enhanced Kit . |Exposure time: 90s.



Western blot analysis of extracts of various cell lines, using Pan Mono-Methyl lysine Rabbit pAb (TA379599) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 3min.