

Product datasheet for **TA379608**

Phospho-Tyrosine Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	ICC/IF, WB
Recommended Dilution:	WB,1:500 - 1:2000 IF,1:50 - 1:200 IP,1:50 - 1:200
Reactivity:	Human, Mouse, Rat, Other (Wide Range)
Modifications:	Pan Phospho-Tyrosine
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	A synthetic peptide corresponding to a sequence containing phosphorylated Y.
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.




[View online »](#)


Product images:



Western blot analysis of extracts of Jurkat cells, using Pan Phospho-Tyrosine pAb (TA379608) at 1:1000 dilution. Jurkat cells were treated by Pervanadate (1 mM) at 37°C for 30 minutes. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA. | Detection: ECL Basic Kit. | Exposure time: 1s.



Immunofluorescence analysis of C6 cells using Pan Phospho-Tyrosine (TA379608) at dilution of 1:100. Blue: DAPI for nuclear staining. C6 cells were treated by 20% FBS at 37°C for 20 minutes after serum-starvation overnight.



Immunofluorescence analysis of NIH/3T3 cells using Pan Phospho-Tyrosine (TA379608) at dilution of 1:100. Blue: DAPI for nuclear staining. NIH/3T3 cells were treated by 20% FBS at 37°C for 20 minutes after serum-starvation overnight.