

Product datasheet for TA379711S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PHOS (PDC) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC-P,1:50 - 1:200

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Mouse, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 28kDa

Gene Name: phosducin **Database Link:** P20941

Background: This gene encodes a phosphoprotein, which is located in the outer and inner segments of the

rod cells in the retina. This protein may participate in the regulation of visual

phototransduction or in the integration of photoreceptor metabolism. It modulates the phototransduction cascade by interacting with the beta and gamma subunits of the retinal G-protein transducin. This gene is a potential candidate gene for retinitis pigmentosa and Usher syndrome type II. Alternatively spliced transcript variants encoding different isoforms have


been identified.

Synonyms: MEKA; PHD; PhLOP; PhLP; phosducin






Product images:



Western blot analysis of various lysates using PDC Rabbit pAb ([TA379711]) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit lgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Immunohistochemistry analysis of paraffinembedded Rat retina using PDC Rabbit pAb ([TA379711]) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.