

Product datasheet for TA379746

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PDGF Receptor alpha (PDGFRA) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB,1:500 - 1:2000

Reactivity: Human

Modifications: Phospho Y849

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A synthetic phosphorylated peptide around Y849 of human PDGFRA (NP_006197.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 24kDa/82kDa/122kDa

Gene Name: platelet derived growth factor receptor alpha

Database Link: Entrez Gene 5156 Human

P16234

Background: This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived

growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The

identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived growth factor receptor alpha and beta polypeptides. Studies suggest that this gene plays a role in organ development, wound healing, and tumor progression. Mutations in this gene have been associated with idiopathic hypereosinophilic syndrome, somatic and familial

gastrointestinal stromal tumors, and a variety of other cancers.






Synonyms:

CD140A; MGC74795; PDGF-R-alpha; PDGFR2; RHEPDGFRA

Product images:



Western blot analysis of extracts of HepG2 cells, using Phospho-PDGFRA-Y849 antibody (TA379746) at 1:1000 dilution. HepG2 cells were treated by EGF (100ng/mL) for 30 minutes after serum-starvation overnight. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% BSA.