

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA379767

PDK3 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
lsotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 137-406 of human PDK3 (NP_005382.1).
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	46kDa/48kDa
Gene Name:	pyruvate dehydrogenase kinase 3
Database Link:	<u>Entrez Gene 5165 Human</u> <u>Q15120</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


CRIGENE PDK3 Rabbit Polyclonal Antibody – TA379767

PDK3

Background:The pyruvate dehydrogenase (PDH) complex is a nuclear-encoded mitochondrial
multienzyme complex that catalyzes the overall conversion of pyruvate to acetyl-CoA and
CO(2). It provides the primary link between glycolysis and the tricarboxylic acid (TCA) cycle,
and thus is one of the major enzymes responsible for the regulation of glucose metabolism.
The enzymatic activity of PDH is regulated by a phosphorylation/dephosphorylation cycle,
and phosphorylation results in inactivation of PDH. The protein encoded by this gene is one
of the three pyruvate dehydrogenase kinases that inhibits the PDH complex by
phosphorylation of the E1 alpha subunit. This gene is predominantly expressed in the heart
and skeletal muscles. Alternatively spliced transcript variants encoding different isoforms
have been found for this gene.

Synonyms:

Product images:



Western blot analysis of extracts of various cell lines, using PDK3 antibody (TA379767) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 1s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US