

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA379869

PHF19 Rabbit Polyclonal Antibody

Product data:

| Product Type: | Primary Antibodies |
|-------------------------|--|
| Applications: | ICC/IF, WB |
| Recommended Dilution: | WB,1:500 - 1:2000 IF,1:50 - 1:100 |
| Reactivity: | Human, Mouse, Rat |
| Modifications: | Unmodified |
| Host: | Rabbit |
| lsotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 1-207 of human PHF19 (NP_001009936.1). |
| Formulation: | PBS with 0.09% Sodium azide,50% glycerol,pH7.3. |
| Concentration: | lot specific |
| Purification: | Affinity purification |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. |
| Stability: | Shelf life: one year from despatch. |
| Predicted Protein Size: | 14kDa/22kDa/65kDa |
| Gene Name: | PHD finger protein 19 |
| Database Link: | <u>Entrez Gene 26147 Human</u> <u>Q5T6S3</u> |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


GRIGENE PHF19 Rabbit Polyclonal Antibody – TA379869

Background:Polycomb group (PcG protein that specifically binds histone H3 trimethylated at 'Lys-36'
(H3K36me3 and recruits the PRC2 complex, thus enhancing PRC2 H3K27me3 methylation
activity. Probably involved in the transition from an active state to a repressed state in
embryonic stem cells: acts by binding to H3K36me3, a mark for transcriptional activation, and
recruiting H3K36me3 histone demethylases RIOX1 or KDM2B, leading to demethylation of
H3K36 and recruitment of the PRC2 complex that mediates H3K27me3 methylation, followed
by de novo silencing. Recruits the PRC2 complex to CpG islands and contributes to embryonic
stem cell self-renewal. Also binds histone H3 dimethylated at 'Lys-36' (H3K36me2. Isoform 1
and isoform 2 inhibit transcription from an HSV-tk promoter.


Synonyms:

DKFZP727G051; hPCL3; MGC23929; MGC131698; MGC149712; MGC149713; MTF2L1; PCL3

Product images:



Western blot analysis of extracts of various cell lines, using PHF19 antibody (TA379869) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 90s.



Immunofluorescence analysis of A-549 cells using PHF19 antibody (TA379869).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US