

Product datasheet for TA379915

FAM38A (PIEZO1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 2230-2420

of human PIEZO1 (NP_001136336.2).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 286kDa

Gene Name: piezo type mechanosensitive ion channel component 1

Database Link: Entrez Gene 9780 Human

Q92508

Background: The protein encoded by this gene is a mechanically-activated ion channel that links

mechanical forces to biological signals. The encoded protein contains 36 transmembrane domains and functions as a homotetramer. Defects in this gene have been associated with

dehydrated hereditary stomatocytosis.

Synonyms: 9630020g22; Fam38a; mKIAA0233




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Product images:



Western blot analysis of extracts of Mouse kidney, using PIEZO1 antibody (TA379915) at 1:1000 dilution.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit.|Exposure time: 90s.



Western blot analysis of extracts of Mouse lung, using PIEZO1 antibody (TA379915) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 90s.



Immunofluorescence analysis of NIH-3T3 cells using PIEZO1 Rabbit pAb (TA379915) at dilution of 1:250 (40x lens). Blue: DAPI for nuclear staining.

