

### **Product datasheet for TA380017**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Phospholipase C gamma 1 (PLCG1) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

**Reactivity:** WB,1:500 - 1:2000 Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A synthetic peptide of human PLC gamma 1 (PLC gamma 1 (PLCG1)).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 148kDa

Gene Name: phospholipase C gamma 1

Database Link: Entrez Gene 5335 Human

<u>P19174</u>

**Background:** The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and

diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript

variants encoding different isoforms have been found for this gene.


**Synonyms:** 1-phosphatidyl-D-myo-inositol-4,5-bisphosphate; inositoltrisphosphohydrolase; NCKAP3;

OTTHUMP00000178982; PLC-148; PLC-gamma-1; PLC-II; PLC1; PLC148; PLCgamma1





# **Product images:**



Western blot analysis of extracts of various cell lines, using PLC gamma 1 (PLC gamma 1 (PLCG1)) Rabbit pAb (TA380017) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.