

Product datasheet for TA380073S

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

Noxa (PMAIP1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, ICC/IF, WB

Recommended Dilution: WB.1:2000 - 1:6000

IF/ICC,1:50 - 1:200

ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 6kDa

Gene Name: phorbol-12-myristate-13-acetate-induced protein 1

Database Link: Entrez Gene 5366 Human

Q13794

Background: This gene belongs to a pro-apoptotic subfamily within the BCL-2 protein family, referred to as

the BCL-2 homology domain 3 (BH3)-only subfamily, which determine whether a cell commits to apoptosis. In response to death-inducing stimuli, BH3-only members inhibit the anti-apoptotic BCL-2 family members, which under steady-state conditions keep the multi-BH


domain proteins BAX and BAK, in an inactive state.

Synonyms: APR; NOXA






Product images:





Western blot analysis of various lysates



Immunofluorescence analysis of Jurkat cells using PMAIP1 Rabbit pAb ([TA380073]) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.







Immunofluorescence analysis of paraffinembedded rat thymus using PMAIP1 Rabbit pAb ([TA380073]) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of paraffinembedded mouse thymus using PMAIP1 Rabbit pAb ([TA380073]) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.