

## Product datasheet for **TA380081**

### PML Protein (PML) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB,1:500 - 1:2000
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant protein of human PML/RARA
Formulation:	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	47-48kDa/62-97kDa/39kDa/50kDa
Gene Name:	promyelocytic leukemia
Database Link:	<a href="#">Entrez Gene 5371 Human P29590</a>




[View online »](#)

**Background:**

Promyelocytic leukemia/retinoic acid receptor alpha or PML-RARA refers to an abnormal fusion gene sequence. It is a specific rearrangement of genetic material from two separate chromosomes (chromosomal translocation) and is associated with a specific type of leukemia. Promyelocytic leukemia (PML) is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This phosphoprotein localizes to nuclear bodies where it functions as a transcription factor and tumor suppressor. Its expression is cell-cycle related and it regulates the p53 response to oncogenic signals. The gene is often involved in the translocation with the retinoic acid receptor alpha gene associated with acute promyelocytic leukemia (APL). Retinoic acid receptor alpha (RARA), regulates transcription in a ligand-dependent manner. This gene has been implicated in regulation of development, differentiation, apoptosis, granulopoiesis, and transcription of clock genes. Translocations between this locus and several other loci have been associated with acute promyelocytic leukemia.

**Synonyms:**

MYL; PP8675; RNF71; TRIM19

**Product images:**

Western blot analysis of extracts of various cells, using PML/RARA antibody (TA380081) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 90s.