

Product datasheet for TA380229

PPP1A (PPP1CA) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Reactivity: WB,1:500 - 1:2000 Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-330 of

human PPP1CA (NP_002699.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 32kDa/37kDa/38kDa

Gene Name: protein phosphatase 1 catalytic subunit alpha

Database Link: Entrez Gene 5499 Human

P62136

Background: The protein encoded by this gene is one of the three catalytic subunits of protein

phosphatase 1 (PP1). PP1 is a serine/threonine specific protein phosphatase known to be involved in the regulation of a variety of cellular processes, such as cell division, glycogen metabolism, muscle contractility, protein synthesis, and HIV-1 viral transcription. Increased PP1 activity has been observed in the end stage of heart failure. Studies in both human and mice suggest that PP1 is an important regulator of cardiac function. Mouse studies also suggest that PP1 functions as a suppressor of learning and memory. Three alternatively spliced transcript variants encoding different isoforms have been found for this gene.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

MGC1674; MGC15877; PP-1A; PP1A; PPP1A

Product images:



Western blot analysis of extracts of various cell lines, using PPP1CA antibody (TA380229) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit .