

Product datasheet for TA380231S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Protein Phosphatase 1 beta (PPP1CB) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200 IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit

Isotype: IgG
Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-327 of

human PP1 beta (NP_002700.1).

Formulation: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 37kDa

Gene Name: protein phosphatase 1 catalytic subunit beta

Database Link: Entrez Gene 5500 Human

P62140

Background: The protein encoded by this gene is one of the three catalytic subunits of protein

phosphatase 1 (PP1). PP1 is a serine/threonine specific protein phosphatase known to be involved in the regulation of a variety of cellular processes, such as cell division, glycogen metabolism, muscle contractility, protein synthesis, and HIV-1 viral transcription. Mouse

studies suggest that PP1 functions as a suppressor of learning and memory. Two

alternatively spliced transcript variants encoding distinct isoforms have been observed.







Synonyms:

MGC3672; OTTHUMP00000200968; PP-1B; PPP1CD

Product images:





Western blot analysis of extracts of various cell lines, using PP1 beta antibody ([TA380231]) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit. | Exposure time: 60s.



Immunofluorescence analysis of U2OS cells using PP1 beta Rabbit pAb ([TA380231]) at dilution of 1:100. Blue: DAPI for nuclear staining.







Immunofluorescence analysis of L929 cells using PP1 beta Rabbit pAb ([TA380231]) at dilution of 1:100. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of H9C2 cells using PP1 beta Rabbit pAb ([TA380231]) at dilution of 1:100. Blue: DAPI for nuclear staining.