

## **Product datasheet for TA380288**

## 1104400 4444511000117150020

## **PRB4 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WE

Recommended Dilution: WB,1:500 - 1:2000

**Reactivity:** Mouse, Rat **Modifications:** Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 178-247 of

human PRB4 (NP\_002714.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

**Predicted Protein Size:** 31kDa

**Gene Name:** proline-rich protein BstNI subfamily 4

Database Link: P10163

**Background:** This gene encodes a member of the heterogeneous family of basic, proline-rich, human

salivary glycoproteins. The encoded preproprotein undergoes proteolytic processing to generate one or more mature peptides before secretion from the parotid glands. Multiple alleles of this gene exhibiting variations in the length of the tandem repeats have been identified. The reference genome encodes the "Small" allele. This gene is located in a cluster of closely related salivary proline-rich proteins on chromosome 12. Alternative splicing results

in multiple transcript variants encoding different isoforms that may undergo similar

proteolytic processing.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: Po

## **Product images:**



Western blot analysis of extracts of various cell lines, using PRB4 antibody (TA380288) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 3min.