

## **Product datasheet for TA380303**

## **PRDM9 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

**Reactivity:** WB,1:500 - 1:2000 **Reactivity:** Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

**Clonality:** Polyclonal

**Immunogen:** Recombinant protein of human PRDM9.

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 103kDa

**Gene Name:** PR domain 9

Database Link: Entrez Gene 56979 Human

Q9NQV7

**Background:** The protein encoded by this gene is a zinc finger protein with histone methyltransferase

activity that catalyzes histone H3 lysine 4 trimethylation (H3K4me3) during meiotic prophase. This protein contains multiple domains, including a Kruppel-associated box (KRAB) domain, an SSX repression domain (SSXRD), a PRD1-BF1 and RIZ homologous region, a subclass of SET (PR/SET) domain, and a tandem array of C2H2 zinc fingers. The zinc finger array recognizes a short sequence motif, leading to local H3K4me3, and meiotic recombination hotspot activity. The observed allelic variation alters the DNA-binding sequence specificity of the protein, resulting in distinct meiotic recombination hotspots amongst individuals and populations.

Multiple alternate alleles of this gene have been described.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

PFM6

## **Product images:**



Western blot analysis of extracts of various cell lines, using PRDM9 antibody (TA380303) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 30s.