

Product datasheet for TA380324

AMPK alpha 1 (PRKAA1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies Applications: IHC, WB Recommended Dilution: WB.1:500 - 1:2000 IHC,1:50 - 1:200 **Reactivity:** Human, Mouse, Rat Host: Rabbit lgG Isotype: **Clonality:** Polyclonal Immunogen: A synthetic peptide of human AMPKa1 alpha 1 Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. **Concentration:** lot specific **Purification:** Affinity purification **Conjugation:** Unconjugated Storage: Store at -20°C. Avoid freeze / thaw cycles. Stability: Shelf life: one year from despatch. **Predicted Protein Size:** 64kDa/65kDa Gene Name: protein kinase AMP-activated catalytic subunit alpha 1 Database Link: Entrez Gene 5562 Human Q13131 Background: The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed.

Synonyms:

AMPK; AMPK1; AMPKa1; MGC33776; MGC57364

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


OriGene Technologies, Inc.

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Product images:



Western blot analysis of extracts of various cell lines, using AMPKa1 alpha 1 antibody (TA380324) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Enhanced Kit . | Exposure time: 3min.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US