

# **Product datasheet for TA380386**

## **PRMT1 Rabbit Polyclonal Antibody**

### **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200 IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 20-353 of

human PRMT1 (NP\_938074.2).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 39kDa/40kDa/41kDa

**Gene Name:** protein arginine methyltransferase 1

**Database Link:** Entrez Gene 3276 Human

Q99873



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### Background:


This gene encodes a member of the protein arginine N-methyltransferase (PRMT) family. Post-translational modification of target proteins by PRMTs plays an important regulatory role in many biological processes, whereby PRMTs methylate arginine residues by transferring methyl groups from S-adenosyl-L-methionine to terminal guanidino nitrogen atoms. The encoded protein is a type I PRMT and is responsible for the majority of cellular arginine methylation activity. Increased expression of this gene may play a role in many types of cancer. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 5.

Synonyms: ANM1; HCP1; HMT2; HRMT1L2; IR1B4

### **Product images:**



Western blot analysis of extracts of various cell lines, using PRMT1 antibody (TA380386) at 1:1000 dilution. |Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. |Lysates/proteins: 25ug per lane. |Blocking buffer: 3% nonfat dry milk in TBST. |Detection: ECL Basic Kit. |Exposure time: 1s.



Immunofluorescence analysis of L929 cells using PRMT1 Rabbit pAb (TA380386) at dilution of 1:100. Blue: DAPI for nuclear staining.