

Product datasheet for **TA380483S**


PTCD3 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, ICC/IF, WB
Recommended Dilution:	WB, 1:500 - 1:2000 IF/ICC, 1:50 - 1:200 ELISA, Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Reactivity:	Human, Mouse, Rat
Modifications:	Unmodified
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Formulation:	Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.
Concentration:	lot specific
Purification:	Affinity purification
Conjugation:	Unconjugated
Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Stability:	Shelf life: one year from despatch.
Predicted Protein Size:	79kDa
Gene Name:	pentatricopeptide repeat domain 3
Database Link:	Entrez Gene 55037 Human Q96EY7
Background:	Enables rRNA binding activity and ribosomal small subunit binding activity. Involved in mitochondrial translation. Located in several cellular components, including cytosol; mitochondrion; and nucleoplasm. Implicated in combined oxidative phosphorylation deficiency 51.
Synonyms:	DKFZp666K071; FLJ20758; TRG-15


[View online »](#)

Product images:



Western blot analysis of various lysates using PTCD3 Rabbit pAb ([TA380483]) at 1:1000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using PTCD3 Rabbit pAb ([TA380483]) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.