

Product datasheet for TA380506S

PTGS1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:100 - 1:200 IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 1-180 of

human COX1/COX1/PTGS1 (NP 001258094.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 56kDa/61kDa/64kDa/68kDa/71kDa/72kDa **Gene Name:** prostaglandin-endoperoxide synthase 1

Database Link: Entrez Gene 5742 Human

P23219

Background: This is one of two genes encoding similar enzymes that catalyze the conversion of

arachinodate to prostaglandin. The encoded protein regulates angiogenesis in endothelial cells, and is inhibited by nonsteroidal anti-inflammatory drugs such as aspirin. Based on its ability to function as both a cyclooxygenase and as a peroxidase, the encoded protein has been identified as a moonlighting protein. The protein may promote cell proliferation during

tumor progression. Alternative splicing results in multiple transcript variants.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

COX-1; COX1; COX3; Cyclooxygenase-1; PCOX1; PGG/HS; PGHS-1; PGHS1; PTGHS

Product images:



Western blot analysis of extracts of various cell lines, using COX1/COX1/PTGS1 antibody ([TA380506]) at 1:1000 dilution._Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution._Lysates/proteins: 25ug per lane._Blocking buffer: 3% nonfat dry milk in TBST._Detection: ECL Enhanced Kit ._Exposure time: 15s.



Immunofluorescence analysis of HeLa cells using COX1/COX1/PTGS1 Polyclonal Antibody ([TA380506]) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.