

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA380520

CCK4 (PTK7) Rabbit Polyclonal Antibody


Product data:

Applications:IHC WBApplications:WB 1:500 1:2000 IHC 1:50 1:100Reactivity:Human, Mouse, RatModifications:UnmodifiedMotifications:RabitIsotype:ababtIsotype:OlyclonalClonality:PolyclonalImmunogen:Buffer: PBS with 0.09% Sodium azide, 50% glycerol, PH7.3.Formulation:NonoligatedPorrigitation:Atfinity purificationPorrigitation:Store at-20°C. Avoid freeze / thaw cycles.Storage:Store at-20°C. Avoid freeze / thaw cycles.Storage:Store at-20°C. Avoid freeze / thaw cycles.Predicted Proteinsize:Resonal fusion Protein containing a sequence corresponding to amino acids Resonant fusion protein containing a sequence corresponding to amino acids Resonant fusion protein containing a sequence corresponding to amino acids Resonant fusion protein containing a sequence corresponding to amino acids Resonant fusion protein containing a sequence corresponding to amino acids Resonant fusion protein containing a sequence corresponding to amino acids Resonant fusion protein containing a sequence corresponding to amino acids Resonant fusion protein containing a sequence corresponding to amino acids Resonant fusion protein containing a sequence corresponding to amino acids Resonant fusion protein containing a sequence corresponding to amino acids Resonant fusion protein containing a sequence corresponding to acids Resonant fusion protein contain	Product Type:	Primary Antibodies
IHC,1:50-1:100Reactivity:Human, Mouse, RatModifications:UnmodifiedHost:RabbitIsotype:IgGClonality:PolyclonalImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 811-1070 of human PTK7 (NP_002812.2).Formulation:Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.Concentration:Ids specificPurification:Auffnity purificationKorage:Store at -20°C. Avoid freeze / thaw cycles.Stability:Shelf life: one year from despatch.Predicted Protein SizeBykDa/103kDa/112kDa/113kDa/119kDaGene Name:Entrez Gene 5754 Human Oli3308Background:This gene encodes a member of the receptor protein tyrosine kinase family of protein sthat strasduce extracellular signals across the cell membrane. The encoded protein lacks arei nuntliple cellular processes including polarity and adhesion. Alternatively spliced area curity variants encoding multiple isoforms have been observed for this gene.	Applications:	IHC, WB
ModifiedModifiedHost:RabbitIsotype:IgGClonality:PolyclonalImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 811-1070 of human PTK7 (NP_002812.2).Formulation:Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.Concentration:Iot specificPurification:UnconjugatedStorage:Store at -20°C. Avoid freeze / thaw cycles.Stability:Shell file: one year from despatch.Predicted Protein Size:8yRoJ/103kDa/112kDa/113kDa/119kDa/119kDaGene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human O13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks arole in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Recommended Dilution:	
Hose:RabbitIsotype:IgGClonality:PolyclonalImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 811-1070 of human PTK7 (NP_002812.2).Formulation:Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.Forneration:Iot specificPurification:Affnity purificationPurification:Minon protein containing a sequence corresponding to amino acids 811-1070 of human PTK7 (NP_002812.2).Fornulation:Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.Purification:Not specificForniguetion:Minon protein containing a sequence corresponding to amino acids 811-1070 of human PTK7 (NP_002812.2).Foriguetion:Not specificForiguetion:Not specificStability:Not specificStability:Sole at -20°C. Avoid freeze / thaw cycles.Fredicted Protein SizeSoleAl/103kDa/112kDa/113kDa/119kDaPredicted Protein SizeSoleAl/103kDa/112kDa/113kDa/119kDaOrabase Link:Entrez Gene 5754 Human QU3308Background:This gene encodes a member of the receptor protein tyrosine kinase family of protein backs detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays arole in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Reactivity:	Human, Mouse, Rat
Isotype:IgGClonality:PolyclonalImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 811-1070 of human PTK7 (NP_002812.2).Formulation:Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.Concentration:Iot specificPurification:Affinity purificationConjugation:UnconjugatedStorage:Sotre at -20°C. Avoid freeze / thaw cycles.Stability:Shelf life: one year from despatch.Predicted Protein Size8yBoJr103kDa/112kDa/118kDa/119kDaGene Name:Interz Gene 5754 Human O13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of protein sthat transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Modifications:	Unmodified
Clonality:PolyclonalImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 811-1070 of human PTK7 (NP_002812.2).Formulation:Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.Concentration:Iot specificPurification:Affinity purificationConjugation:UnconjugatedStorage:Store at -20°C. Avoid freeze / thaw cycles.Stability:Shelf life: one year from despatch.Predicted Protein Size:8ykDa/103kDa/112kDa/113kDa/119kDaGene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Host:	Rabbit
Immunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 811-1070 of human PTK7 (NP_002812.2).Formulation:Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.Concentration:Iot specificPurification:Affinity purificationConjugation:UnconjugatedStorage:Store at -20°C. Avoid freeze / thaw cycles.Stability:Shelf life: one year from despatch.Predicted Protein Size:89kDa/103kDa/112kDa/113kDa/119kDaGene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human O13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	lsotype:	lgG
of human PTK7 (NP_002812.2).Formulation:Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.Concentration:lot specificPurification:Affinity purificationConjugation:UnconjugatedStorage:Store at -20°C. Avoid freeze / thaw cycles.Stability:Shelf life: one year from despatch.Predicted Protein Size:89kDa/103kDa/112kDa/113kDa/119kDaGene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Clonality:	Polyclonal
Concentration:lot specificPurification:Affinity purificationConjugation:UnconjugatedStorage:Store at -20°C. Avoid freeze / thaw cycles.Stability:Shelf life: one year from despatch.Predicted Protein Size:89kDa/103kDa/112kDa/113kDa/119kDaGene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Immunogen:	
Purification:Affinity purificationConjugation:UnconjugatedStorage:Store at -20°C. Avoid freeze / thaw cycles.Stability:Shelf life: one year from despatch.Predicted Protein Size:89kDa/103kDa/112kDa/113kDa/119kDaGene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Formulation:	Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.
Conjugation:UnconjugatedStorage:Store at -20°C. Avoid freeze / thaw cycles.Stability:Shelf life: one year from despatch.Predicted Protein Size:89kDa/103kDa/112kDa/113kDa/119kDaGene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Concentration:	lot specific
Storage:Store at -20°C. Avoid freeze / thaw cycles.Stability:Shelf life: one year from despatch.Predicted Protein Size:89kDa/103kDa/112kDa/113kDa/118kDa/119kDaGene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Purification:	Affinity purification
Stability:Shelf life: one year from despatch.Predicted Protein Size:89kDa/103kDa/112kDa/113kDa/118kDa/119kDaGene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Conjugation:	Unconjugated
Predicted Protein Size:89kDa/103kDa/112kDa/113kDa/118kDa/119kDaGene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Storage:	Store at -20°C. Avoid freeze / thaw cycles.
Gene Name:protein tyrosine kinase 7 (inactive)Database Link:Entrez Gene 5754 Human Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Stability:	Shelf life: one year from despatch.
Database Link:Entrez Gene 5754 Human Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Predicted Protein Size:	89kDa/103kDa/112kDa/113kDa/118kDa/119kDa
Q13308Background:This gene encodes a member of the receptor protein tyrosine kinase family of proteins that transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Gene Name:	protein tyrosine kinase 7 (inactive)
transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.	Database Link:	
Synonyms: CCK-4; CCK4	Background:	transduce extracellular signals across the cell membrane. The encoded protein lacks detectable catalytic tyrosine kinase activity, is involved in the Wnt signaling pathway and plays a role in multiple cellular processes including polarity and adhesion. Alternatively spliced
	Synonyms:	ССК-4; ССК4



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Product images:



Western blot analysis of extracts of various cell lines, using PTK7 antibody (TA380520) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 15s.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US