

## **Product datasheet for TA380540S**

## PTP alpha (PTPRA) Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:100 - 1:200

Reactivity: Human, Mouse

Modifications: Unmodified

**Host:** Rabbit

**Isotype:** IgG

Clonality: Polyclonal

**Immunogen:** Recombinant fusion protein containing a sequence corresponding to amino acids 20-142 of

human PTPRA (NP\_543031.1).

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

**Storage:** Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 89kDa/90kDa

**Gene Name:** protein tyrosine phosphatase, receptor type A

Database Link: Entrez Gene 5786 Human

P18433



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic catalytic domains, and thus represents a receptor-type PTP. This PTP has been shown to dephosphorylate and activate Src family tyrosine kinases, and is implicated in the regulation of integrin signaling, cell adhesion and proliferation. Three alternatively spliced variants of this gene, which encode two distinct isoforms, have been reported.

Synonyms:

HEPTP; HLPR; HPTPA; HPTPalpha; LRP; OTTHUMP00000030064; PTPA; PTPase-alpha; PTPRL2; R-PTP-alpha; RPTPA

## **Product images:**



Western blot analysis of extracts of various cell lines, using PTPRA antibody ([TA380540]) at 1:1000 dilution.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Enhanced Kit.|Exposure time: 90s.