

## **Product datasheet for TA380546**

## Froduct datasifeet for TA380340

## **LAR (PTPRF) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: WB,1:1000 - 1:5000

ELISA,Recommended starting concentration is 1 μg/mL. Please optimize the concentration

based on your specific assay requirements.

Reactivity: Human, Mouse

Modifications: Unmodified

**Host:** Rabbit

Isotype: IgG

Clonality: Polyclonal

**Formulation:** Buffer: PBS with 0.09% Sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 213kDa

**Gene Name:** protein tyrosine phosphatase, receptor type F

Database Link: Entrez Gene 5792 Human

<u>P10586</u>



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP possesses an extracellular region, a single transmembrane region, and two tandem intracytoplasmic catalytic domains, and thus represents a receptor-type PTP. The extracellular region contains three Ig-like domains, and nine non-Ig like domains similar to that of neural-cell adhesion molecule. This PTP was shown to function in the regulation of epithelial cell-cell contacts at adherents junctions, as well as in the control of beta-catenin signaling. An increased expression level of this protein was found in the insulin-responsive tissue of obese, insulin-resistant individuals, and may contribute to the pathogenesis of insulin resistance. Two alternatively spliced transcript variants of this gene, which encode distinct proteins, have been reported.

Synonyms:

FLJ43335; FLJ45062; FLJ45567; LAR; LCA-homolog; OTTHUMP00000008683; OTTHUMP00000008684

## **Product images:**



Western blot analysis of lysates from bewo cells