

## Product datasheet for TA380554

## **GLEPP1 (PTPRO) Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** WB

Recommended Dilution: WB.1:500 - 1:2000

Reactivity: Mouse

**Modifications:** Unmodified

Host: Rabbit Isotype: **IgG** 

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 30-300 of

human PTPRO (NP\_109592.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

**Predicted Protein Size:** 43kDa/47kDa/67kDa/134kDa/138kDa

Gene Name: protein tyrosine phosphatase, receptor type O

**Database Link:** Q16827

Background: This gene encodes a member of the R3 subtype family of receptor-type protein tyrosine

> phosphatases. These proteins are localized to the apical surface of polarized cells and may have tissue-specific functions through activation of Src family kinases. This gene contains two distinct promoters, and alternatively spliced transcript variants encoding multiple isoforms have been observed. The encoded proteins may have multiple isoform-specific and tissuespecific functions, including the regulation of osteoclast production and activity, inhibition of cell proliferation and facilitation of apoptosis. This gene is a candidate tumor suppressor, and

decreased expression of this gene has been observed in several types of cancer.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms: GLEPP1; PTP-oc; PTP-U2; PTPROt; PTPU2

## **Product images:**



Western blot analysis of extracts of HT-29 cells, using PTPRO Antibody (TA380554) at 1:1000 dilution.\_Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.\_Lysates/proteins: 25ug per lane.\_Blocking buffer: 3% nonfat dry milk in TBST.\_Detection: ECL Enhanced Kit .\_Exposure time: 30s.