

Product datasheet for TA380665

RAD18 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ICC/IF, WB

Recommended Dilution: WB,1:500 - 1:2000

IHC,1:50 - 1:200 IF,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 216-495 of

human Rad18 (NP_064550.3).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 56kDa

Gene Name: RAD18, E3 ubiquitin protein ligase

Database Link: <u>Entrez Gene 56852 Human</u>

O9NS91

Background: The protein encoded by this gene is highly similar to S. cerevisiae DNA damage repair protein

Rad18. Yeast Rad18 functions through its interaction with Rad6, which is an ubiquitin-

conjugating enzyme required for post-replication repair of damaged DNA. Similar to its yeast counterpart, this protein is able to interact with the human homolog of yeast Rad6 protein through a conserved ring-finger motif. Mutation of this motif results in defective replication

of UV-damaged DNA and hypersensitivity to multiple mutagens.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

hHR18; hRAD18; RNF73

Product images:




Western blot analysis of extracts of various cell lines, using Rad18 antibody (TA380665) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 120s.



Immunofluorescence analysis of U2OS cells using Rad18 antibody (TA380665). Blue: DAPI for nuclear staining.





Immunofluorescence analysis of U2OS cells using Rad18 antibody (TA380665). Blue: DAPI for nuclear staining. DNA damage by a UV-A laser.