

# **Product datasheet for TA380765**

# Rb2 p130 (RBL2) Rabbit Polyclonal Antibody

## **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type:           | Primary Antibodies  |
|-------------------------|---|
| Applications:           | ICC/IF, WB  |
| Recommended Dilution:   | WB,1:500 - 1:2000<br>IF,1:50 - 1:200  |
| Reactivity:             | Human, Mouse  |
| Modifications:          | Unmodified  |
| Host:                   | Rabbit  |
| lsotype:                | lgG   |
| Clonality:              | Polyclonal  |
| Immunogen:              | A synthetic peptide corresponding to a sequence within amino acids 100-200 of human RBL2 (NP_005602.3). |
| Formulation:            | Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.   |
| Concentration:          | lot specific  |
| Purification:           | Affinity purification   |
| Conjugation:            | Unconjugated  |
| Storage:                | Store at -20°C. Avoid freeze / thaw cycles.   |
| Stability:              | Shelf life: one year from despatch.   |
| Predicted Protein Size: | 59kDa/128kDa  |
| Gene Name:              | RB transcriptional corepressor like 2   |
| Database Link:          | <u>Entrez Gene 5934 Human</u><br><u>Q08999</u>  |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


#### **GRIGENE** Rb2 p130 (RBL2) Rabbit Polyclonal Antibody – TA380765

Background:Key regulator of entry into cell division. Directly involved in heterochromatin formation by<br/>maintaining overall chromatin structure and, in particular, that of constitutive<br/>heterochromatin by stabilizing histone methylation. Recruits and targets histone<br/>methyltransferases KMT5B and KMT5C, leading to epigenetic transcriptional repression.<br/>Controls histone H4 'Lys-20' trimethylation. Probably acts as a transcription repressor by<br/>recruiting chromatin-modifying enzymes to promoters. Potent inhibitor of E2F-mediated<br/>trans-activation, associates preferentially with E2F5. Binds to cyclins A and E. Binds to and<br/>may be involved in the transforming capacity of the adenovirus E1A protein. May act as a<br/>tumor suppressor.


Synonyms:

FLJ26459; p130; PRB2; Rb2; RBR-2

### **Product images:**



Western blot analysis of extracts of Neuro-2a cells, using RBL2 antibody (TA380765) at 1:1000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit . | Exposure time: 120s.



Immunofluorescence analysis of U2OS cells using RBL2 Rabbit pAb (TA380765) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US