

## **Product datasheet for TA380941**

## Troduct datastreet for TASOUS4

## **Ripk3 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

Applications: WB

**Reactivity:** WB,1:500 - 1:2000 Human, Mouse

Modifications: Phospho T231/S232

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** A phospho specific peptide corresponding to residues surrounding T231/S232 of Mouse

Ripk3.

**Formulation:** Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

**Concentration:** lot specific

**Purification:** Affinity purification

**Conjugation:** Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

**Stability:** Shelf life: one year from despatch.

Predicted Protein Size: 53kDa

**Gene Name:** receptor-interacting serine-threonine kinase 3

Database Link: Entrez Gene 56532 Mouse

Q9QZL0

**Background:** The product of this gene is a member of the receptor-interacting protein (RIP) family of

serine/threonine protein kinases, and contains a C-terminal domain unique from other RIP family members. The encoded protein is predominantly localized to the cytoplasm, and can undergo nucleocytoplasmic shuttling dependent on novel nuclear localization and export signals. It is a component of the tumor necrosis factor (TNF) receptor-I signaling complex,

and can induce apoptosis and weakly activate the NF-kappaB transcription factor.

Synonyms: RIP-3; RIP3




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Western blot analysis of extracts of L929 cells, using Phospho-RIP3-T231/S232 antibody (TA380941) at 1:500 dilution.L929 cells were treated by Calyculin A (100 nM) at 37°C for 30 minutes after serum-starvation overnight. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 180s.