

Product datasheet for TA381138

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

S6K1 (RPS6KB1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Reactivity: WB,1:500 - 1:2000 Human, Mouse

Modifications: Phospho-RPS6KB1-S371

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: A phospho specific peptide corresponding to residues surrounding S371 of human RPS6KB1.

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 51kDa/52kDa/56kDa/59kDa **Gene Name:** ribosomal protein S6 kinase B1

Database Link: Entrez Gene 6198 Human

P23443

Background: This gene encodes a member of the ribosomal S6 kinase family of serine/threonine kinases.

The encoded protein responds to mTOR (mammalian target of rapamycin) signaling to promote protein synthesis, cell growth, and cell proliferation. Activity of this gene has been associated with human cancer. Alternatively spliced transcript variants have been observed. The use of alternative translation start sites results in isoforms with longer or shorter N-termini which may differ in their subcellular localizations. There are two pseudogenes for this


gene on chromosome 17. [provided by RefSeq, Jan 2013]

Synonyms: p70(S6K)-alpha; p70-alpha; p70-S6K; P70S6K; PS6K; S6K1; STK14A





Product images:



Western blot analysis of extracts of NIH/3T3, using Phospho-p70 S6 Kinase 1-S371 antibody (TA381138) at 1:1000 dilution.NIH/3T3 cells were treated by 10% FBS at 37°C for 30 minutes after serum-starvation overnight.NIH/3T3 cells were treated by EGF (100 ng/ml) at 37°C for 30 minutes after serum-starvation overnight.|Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit.|Exposure time: 30s.