

Product datasheet for TA381192

RUNX1T1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:200 - 1:1000

Reactivity: Human

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 100-390 of

human RUNX1T1 (NP_783554.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 44kDa/48kDa/63kDa/64kDa/67kDa/68kDa

Gene Name: RUNX1 translocation partner 1

Database Link: Entrez Gene 862 Human

Q06455

Background: This gene encodes a member of the myeloid translocation gene family which interact with

DNA-bound transcription factors and recruit a range of corepressors to facilitate

transcriptional repression. The t(8;21)(q22;q22) translocation is one of the most frequent karyotypic abnormalities in acute myeloid leukemia. The translocation produces a chimeric gene made up of the 5'-region of the runt-related transcription factor 1 gene fused to the 3'-

region of this gene. The chimeric protein is thought to associate with the nuclear

corepressor/histone deacetylase complex to block hematopoietic differentiation. Alternative

splicing results in multiple transcript variants.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Synonyms:

AML1T1; CBFA2T1; CDR; ETO; MGC2796; MTG8; MTG8b; ZMYND2

Product images:



Western blot analysis of extracts of various cell lines, using RUNX1T1 antibody (TA381192) at 1:1000 dilution.|Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution.|Lysates/proteins: 25ug per lane.|Blocking buffer: 3% nonfat dry milk in TBST.|Detection: ECL Basic Kit.|Exposure time: 1s.