

Product datasheet for TA381352

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Semaphorin 7a (SEMA7A) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB,1:500 - 1:1000

Reactivity: Human

Modifications: Unmodified

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 45-280 of

human SEMA7A (NP_003603.1).

Formulation: Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Concentration: lot specific

Purification: Affinity purification

Conjugation: Unconjugated

Storage: Store at -20°C. Avoid freeze / thaw cycles.

Stability: Shelf life: one year from despatch.

Predicted Protein Size: 73kDa/74kDa

Gene Name: semaphorin 7A (John Milton Hagen blood group)

Database Link: Entrez Gene 8482 Human

<u>075326</u>

Background: This gene encodes a member of the semaphorin family of proteins. The encoded

preproprotein is proteolytically processed to generate the mature

glycosylphosphatidylinositol (GPI)-anchored membrane glycoprotein. The encoded protein is

found on activated lymphocytes and erythrocytes and may be involved in

immunomodulatory and neuronal processes. The encoded protein carries the John Milton Hagen (JMH) blood group antigens. Mutations in this gene may be associated with reduced bone mineral density (BMD). Alternative splicing results in multiple transcript variants, at least

one of which encodes an isoform that is proteolytically processed.






Synonyms:

CD108; CDw108; H-SEMA-K1; H-Sema-L; JMH; MGC126692; MGC126696; SEMAK1; SEMAL; Semaphorin-K1; Semaphorin-L

Product images:



Western blot analysis of extracts of various cell lines, using SEMA7A antibody (TA381352) at 1:3000 dilution. | Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. | Lysates/proteins: 25ug per lane. | Blocking buffer: 3% nonfat dry milk in TBST. | Detection: ECL Basic Kit. | Exposure time: 90s.